

Durham Research Online

Deposited in DRO:

05 July 2018

Version of attached file:

Accepted Version

Peer-review status of attached file:

Peer-reviewed

Citation for published item:

Williams, K.S. and Williams, S.T. and Fitzgerald, L.E. and Sheppard, E.C. and Hill, R.A. (2018) 'Brown hyaena and leopard diets on private land in the Soutpansberg Mountains, South Africa.', *African journal of ecology*, 56 (4). pp. 1021-1027.

Further information on publisher's website:

<https://doi.org/10.1111/aje.12539>

Publisher's copyright statement:

This is the accepted version of the following article: Williams, K.S., Williams, S.T., Fitzgerald, L.E., Sheppard, E.C. Hill, R.A. (2018). Brown hyaena and leopard diets on private land in the Soutpansberg Mountains, South Africa. *African Journal of Ecology* 56(4): 1021-1027, which has been published in final form at <https://doi.org/10.1111/aje.12539>. This article may be used for non-commercial purposes in accordance With Wiley Terms and Conditions for self-archiving.

Additional information:

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

**Brown hyaena and leopard diets on private land in the Soutpansberg
Mountains, South Africa**

Kathryn S. Williams ^{a,b*}, Samuel T. Williams ^{a,b,c,d}, Leanne E. Fitzgerald ^{a,b}, Eleanor C.
Sheppard ^b, and Russell A. Hill ^{a,b,c}

^a Department of Anthropology, Durham University, Dawson Building, South Road, Durham,
DH1 3LE, United Kingdom

^b Primate and Predator Project, PO Box 522, Louis Trichardt, 0920, South Africa

^c Department of Zoology, University of Venda, Thohoyandou 0950, South Africa

^d Institute for Globally Distributed Open Research and Education (IGDORE)

*Corresponding author: Kathryn S. Williams (k.s.williams@durham.ac.uk), +27 (0)
710559557

Kathryn S. Williams ORCID id: orcid.org/0000-0002-6890-2196

Samuel T. Williams ORCID id: orcid.org/0000-0001-7205-7426

Russell A. Hill ORCID id: orcid.org/0000-0002-7601-5802

Acknowledgments

We are grateful to the landowners and managers for their participation and granting access to
their properties. We would like to thank Ian Gaigher, Oldrich van Schalkwyk, Caroline
Kruger, Jannie Moolman, Andrea von Gunten, Hannah Birrell, Philippa Goff, Liam Thomas,

Ryan Scott, Carolyn Dunford, and the volunteers from the Earthwatch Institute for their help with data collection and processing. Durham University, the Earthwatch Institute, and an anonymous donor provided funding.

Introduction

Private land comprises a large proportion of the brown hyaena (*Hyaena brunnea*) and leopard (*Panthera pardus*) range, and is vital to their survival (Jacobson et al., 2016; Kent & Hill, 2013). Although prey availability is often highest in protected areas, private land used for game and livestock farming also hosts an abundance of wild and domestic prey (Balme, Slotow, & Hunter, 2010; Kinnaird & O'Brien, 2012).

Scavenging accounts for approximately 95% of the brown hyaena's dietary intake (Maude & Mills, 2005; Mills, 1984; Owens & Owens, 1978). Brown hyaenas depend on large carnivores such as the leopard to kill larger prey species (Mills, 2015; Slater & Muller, 2014; Stein, Fuller, & Marker, 2013). After feeding, leopards often become satiated before they can completely consume large prey animals, so they leave and return to the kill on subsequent occasions to feed further, providing ample scavenging opportunities (Karanth & Sunquist, 2000; Stein et al., 2013).

Studies that compare brown hyaena diets in areas where large predators are either present or absent show significant variation in patterns of consumption and food acquisition between these regions (van der Merwe et al., 2009; Yarnell et al., 2013). The degree of dietary overlap

between brown hyaenas and leopards has, however, rarely been assessed, and this has never been studied in a montane area. We assessed the dietary composition of brown hyaenas and leopards and the degree of dietary overlap between these species in the Soutpansberg Mountains, South Africa. We also compared dietary composition with the relative abundance of prey species.

Methods

Study site

Data were collected from private properties in and around the Soutpansberg Mountains, Limpopo Province, South Africa (Fig. 1). The Soutpansberg Mountains range in altitude from 200 m to 1,748 m above sea level (Berger et al., 2003). Rainfall in the Soutpansberg Mountains ranges from 367 mm to over 2,000 mm per annum (Kabanda, 2003).

Variable climatic conditions and the mountains' undulating topography produce a myriad of biomes which host an extremely high level of biodiversity (Macdonald, Gaigher, Gaigher, & Berger, 2003). The most abundant prey species for large predators in the western Soutpansberg Mountains are bushbuck (*Tragelaphus scriptus*), Cape porcupine (*Hystrix africaeaustralis*), chacma baboon (*Papio ursinus*), greater kudu (*Tragelaphus strepsiceros*), and giraffe (*Giraffa camelopardalis*) (Chase Grey, Bell, & Hill, 2017). Much of the land in the mountains is unsuitable for farming and is used for leisure or ecotourism. Nearby lower lying areas are mainly used for livestock, game, and agricultural farming.

Within the mountains, leopards and brown hyaenas are the only resident large carnivores (Knott, Knott, Kruger, & Van der Waal, 2003). The leopard population in the western Soutpansberg Mountains is suffering a significant population decline (Williams, Williams, Lewis, & Hill, 2017), from 10.7 leopards per 100 km² in 2008 (Chase Grey, Kent, & Hill, 2013) to 3.7 per 100 km² in 2015 (Williams et al., 2017). Illegal human activity is driving high levels of leopard mortality (Williams et al., 2017). In 2015, brown hyaena density was estimated at 3.6 per 100 km² (Williams, 2017). Spotted hyaenas (*Crocuta crocuta*), cheetahs (*Acinonyx jubatus*), African wild dogs (*Lycaon pictus*), and black-backed jackals (*Canis mesomelas*) pass through the area occasionally.

Dietary analysis

Scats were collected opportunistically in the western Soutpansberg Mountains from wild brown hyaenas (n = 137 scats) and leopards (n = 237 scats) between July 1, 2011 and December 31, 2015. Careful consideration of identifying features such as colouration, size, and weight was employed to ensure that scats were correctly assigned to species (Stuart & Stuart, 2003). Since there were no other large carnivores resident, confusion of scats from brown hyaenas and leopards with those from other species was unlikely.

Scats were placed in a wire sieve with 1 mm sized mesh and washed in water to remove all faecal matter (Kuhn, Wiesel, & Skinner, 2008). The contents of the scats were dried in the sun, then spread across a random sampling tray consisting of 36 or 100 numbered squares based on the size of the contents (Martins, Horsnell, Titus, Rautenbach, & Harris, 2011), and the macroscopic qualities of the contents were noted. For all brown hyaena scats and for 75

leopard scats, 40 hairs from every scat were selected at random: 20 hairs were used to create cuticular scale imprints (following Keogh, 1983) and 20 hairs were embedded in clear wax and cross-sectioned (following Douglas, 1989). For the remaining 162 leopard scats, cross-sectional analysis only was conducted. Cuticular imprints and cross-sections were carefully examined under a standard light microscope at 40-100x magnification. The species from which hairs originated were identified by comparing samples with a reference library of hairs collected from known mammal species and with published guides (Keogh, 1983; Seiler, 2010; Taru & Backwell, 2013), and we checked all species identifications at least twice to ensure accuracy.

Camera trapping

An array of 23 camera trap stations composed of two camera traps per station (Reconyx Hyperfire™ HC500 and HC600) was established in the western Soutpansberg Mountains (Fig. 1). The location and spacing of camera stations was optimised for estimation of leopard population density using a spatially explicit capture recapture framework (Williams et al., 2017). All camera stations collected photographs continuously from January 1, 2012 to December 31, 2015. The camera trap array covered an area of 73 km² at the onset of the study, but following one landowner's withdrawal from the survey and the subsequent relocation of five camera stations, the study area was reduced to 59 km² in late 2013 (Williams et al., 2017).

Statistical analysis

All occurrences of a prey item within a scat were calculated as a corrected frequency of occurrence (CFO) (Braczkowski, Watson, Coulson, & Randall, 2012; Henschel, Abernethy, & White, 2005). Employing the CFO accounted for occasions when more than one prey item was detected in a scat. For example, if two species were present in one scat, each species occurrence was weighted at 0.5 (Henschel et al., 2005; Karanth & Sunquist, 1995).

Dietary overlap between brown hyaena and leopard was calculated using Pianka's index (Pianka, 1973):

$$\alpha = \frac{\sum P_{ia}P_{ib}}{\sqrt{\sum P_{ia}^2 \sum P_{ib}^2}}$$

where α equals the dietary overlap between species a and species b , P_{ia} is corrected frequency of occurrence for species a , and P_{ib} is corrected frequency of occurrence for species b . Results range from 0 (no overlap) to 1 (complete overlap) (Pianka, 1973) and values greater than 0.6 were deemed biologically significant (Navia, Mejía-Falla, & Giraldo, 2007). The relationship between the CFO of species in leopard scats and the CFO of species in brown hyaena scats was further tested using linear regression.

Camera trap data were used to estimate the relative abundance of potential prey species consumed by brown hyaena and leopard. Species abundance was calculated using a relative abundance index (RAI) (Negrões et al., 2010; O'Brien, Kinnaird, & Wibisono, 2003):

$$RAI_i = \left(\frac{\sum_j P_{ij}}{\sum_j tn_j} \right) * 100$$

where P_{ij} is the number of independent captures for i th species at j th camera trap location, and tn_j is the total trap-days at the j th camera trap location (Li, McShea, Wang, Shao, & Shi, 2010; O'Brien et al., 2003). Photographs of the same species taken at the same camera station occurring within a 60-minute interval were grouped as a single capture event (Negrões et al., 2010; Rovero & Marshall, 2009), and we excluded species < 1 kg since these are likely to be significantly underrepresented on camera images (Braczkowski et al., 2012; Henschel, Hunter, Coad, Abernethy, & Mühlenberg, 2011).

To determine the relationship between diet composition and prey abundance we used linear regression to test for associations between the RAI and the CFO of prey species in leopard and brown hyaena scats. We excluded Cape porcupine as it was an outlier due to its dense quills defending it from predators (Mori, Maggini, & Menchetti, 2014). Exclusion of this prey species revealed no further obvious influential cases, nor significant deviations from the assumptions of normality and homogeneity of residuals (Quinn & Keough, 2002). All statistical analyses were conducted in R v. 3.3.1 (R Development Core Team, 2017).

Results

Thirty-nine species of mammals were identified in brown hyaena scats and 24 species of mammals were detected in leopard scats (Table 1). Medium sized mammals was the category most frequently consumed by both species (38.80% of the brown hyaena diet and 50.49% in the leopard diet). The five most frequently consumed species by brown hyaena were bushbuck, chacma baboon, common duiker (*Sylvicapra grimmia*), common warthog (*Phacochoerus africanus*), and red duiker (*Cephalophus natalensis*). Three of these species

(bushbuck, chacma baboon, and common duiker) also ranked highest in the leopard diet. Bushbuck was the most commonly consumed prey item for both brown hyaenas and leopards. Livestock (cows (*Bos taurus*), goats (*Capra aegagrus hircus*), and sheep (*Ovis aries*)) accounted for 7.23% of brown hyaena dietary occurrences. No livestock remains were detected in leopard scats. Dietary overlap between leopard and brown hyaena diet was biologically significant, with a Pianka's index of 0.817.

There was a significant positive association between the CFO of prey species in leopard scats and in brown hyaena scats (linear regression: $R^2 = 0.634$, $F_{(1,44)} = 76.17$, $p < 0.001$, gradient = 0.439, intercept = 1.22; Fig. 2a). The relationship remained significant when a potential outlier, bushbuck, was excluded (linear regression: $R^2 = 0.449$, $F_{(1,43)} = 35.02$, $p < 0.001$, gradient = 0.712, intercept = 0.11).

There was a significant positive relationship between the RAI of prey species and their CFO in the scats of brown hyaenas ($R^2 = 0.335$, $F_{(1,24)} = 12.07$, $p = 0.002$, gradient = 0.011, intercept = 2.69; Fig. 2b). No significant relationship was found between the RAI of prey species and their CFO in leopard scats ($R^2 = 0.128$, $F_{(1,15)} = 2.21$, $p = 0.158$).

Discussion

We found that there is high dietary overlap between leopards and brown hyaenas in the Soutpansberg Mountains. Although scat analysis does not definitively explain how prey remains are acquired (Mills & Mills, 1978; Nilsen et al., 2012), our findings support the hypothesis that hyaenas may be acquiring carcasses from leopard kills.

175

176 Scavenging from an apex predator is primarily expected for medium- and large-bodied prey
177 which are unlikely to be completely consumed by leopards immediately after making the kill
178 (Stein et al., 2013; Yarnell et al., 2013), especially since successful brown hyaena hunts are
179 mostly restricted to small- and very small-bodied species (Maude & Mills, 2005). The three
180 most common species in the diets of brown hyaena and leopard are predominantly diurnal
181 (bushbuck 67% diurnal; chacma baboon 99% diurnal; common duiker 78% diurnal:
182 (Fitzgerald, 2015)) with warthogs and red duiker, the fourth and fifth most frequently
183 consumed species by brown hyaena, both 94% diurnal in this area (Fitzgerald, 2015). These
184 species would not be easily accessible to brown hyaenas when hunting, as brown hyaenas
185 have a very low degree of activity during the day (Mills, 1984), so the most likely source of
186 these species is scavenging. In contrast, 36% of leopard activity is during daylight in the
187 Soutpansberg Mountains (Fitzgerald, 2015), and leopards have been observed to hunt diurnal
188 prey at these times. Although it is possible that brown hyaenas have successful hunts during
189 times when prey species are inactive, it is more likely that these species are scavenged.

190

191 Scavenging the remains of animals that died from anthropogenic causes, as well as those that
192 died naturally will also contribute to the brown hyaena's diet. The positive relationship
193 between brown hyaena diet and prey abundance indicates a generalist diet that is common in
194 scavengers (Maude & Mills, 2005). Leopards strongly prefer specific prey species weighing
195 between 10 and 40 kg such as bushbuck (Hayward et al., 2006), which explains why we
196 found no relationship between leopard diet and prey abundance, since very large and small
197 species are not taken in relation to their abundance. Although brown hyaenas may acquire
198 some of their food by scavenging from other predators such as caracals (*Caracal caracal*),

these occur infrequently in the western Soutpansberg Mountains (unpublished data). Nevertheless, leopards appear to represent the greatest opportunity for scavenging in these mountains.

Of the species comprising the brown hyaena diet wild mammals predominated, but a low incidence of feeding on domestic livestock was also noted (7.23% of occurrences). Livestock depredation by brown hyaenas is rare; despite the presence of livestock in the diet of collared brown hyaenas in Botswana, they were never observed hunting livestock (Maude & Mills, 2005). No livestock remains were detected in leopard scats. Yet, leopards do occasionally attack livestock in the area (unpublished data). Therefore, it is likely that brown hyaenas may have secured some livestock remains from leopards residing at lower altitudes or by scavenging the remains of livestock that died from other causes such as disease, mismanagement, or roadkill.

Understanding diet, food acquisition, and interrelationships between predators that are exceedingly reliant on private land is crucial for their conservation. Our data show that brown hyaenas on private land in the Soutpansberg Mountains have a varied diet consisting of mostly wild mammals. High dietary overlap with leopards and evidence supporting scavenging behaviour suggests that leopards could potentially provide brown hyaenas with scavenging opportunities, and thus function as a keystone species for brown hyaenas on private land. Leopards are experiencing severe declines, both in the Soutpansberg Mountains (Williams et al., 2017) and globally (Jacobson et al., 2016). Conservation management plans that adopt a multi-species approach are required to preserve leopards and consequently

provide food security for scavengers like brown hyaenas, which supply important ecosystem services through their feeding habits (Beasley, Olson, & DeVault, 2015).

References

Balme, G. A., Slotow, R., & Hunter, L. T. B. (2010). Edge effects and the impact of non-protected areas in carnivore conservation: leopards in the Phinda-Mkhuze Complex, South Africa. *Animal Conservation*, 13, 315-323. doi:10.1111/j.1469-1795.2009.00342.x

Beasley, J. C., Olson, Z. H., & DeVault, T. L. (2015). Ecological role of vertebrate scavengers. In M. E. Benbow, J. K. Tomberlin, & A. M. Tarone (Eds.), *Carrion Ecology, Evolution, and Their Applications* (pp. 107-121). Boca Raton: CRC Press.

Berger, K., Crafford, J. E., Gaigher, I., Gaigher, M. J., Hahn, N., & Macdonald, I. (2003). Introduction. In K. Berger, J. E. Crafford, I. Gaigher, M. J. Gaigher, N. Hahn, & I. Macdonald (Eds.), *A first synthesis of the environmental, biological and cultural assets of the Soutpansberg* (pp. 1). Louis Trichardt: Leach Printers & Signs.

Brackowski, A., Watson, L., Coulson, D., & Randall, R. (2012). Diet of leopards in the southern Cape, South Africa. *African Journal of Ecology*, 50, 377-380. doi:10.1111/j.1365-2028.2012.01321.x

Chase Grey, J. N., Bell, S., & Hill, R. A. (2017). Leopard diets and landowner perceptions of human wildlife conflict in the Soutpansberg Mountains, South Africa. *Journal for Nature Conservation*, 37, 56-65. doi:10.1016/j.jnc.2017.03.002

242 Chase Grey, J. N., Kent, V. T., & Hill, R. A. (2013). Evidence of a high density population of
 243 harvested leopards in a montane environment. *PLoS ONE*, 8, e82832.
 244 doi:10.1371/journal.pone.0082832

245 Douglas, R. M. (1989). A new method of cross-sectioning hair of larger mammals. *South*
 246 *African Journal of Wildlife Research*, 19, 73-76.

247 Fitzgerald, L. E. (2015). *Diet, activity and spatial occupancy of leopards in the Soutpansberg*
 248 *Mountains*. (MSc), Durham University, Durham University.

249 Hayward, M. W., Henschel, P., O'Brien, J., Hofmeyr, M., Balme, G., & Kerley, G. I. H.
 250 (2006). Prey preferences of the leopard (*Panthera pardus*). *Journal of Zoology*, 270, 298-313.
 251 doi:10.1111/j.1469-7998.2006.00139.x

252 Henschel, P., Abernethy, K. A., & White, L. J. T. (2005). Leopard food habits in the Lopé
 253 National Park, Gabon, Central Africa. *African Journal of Ecology*, 43, 21-28.
 254 doi:10.1111/j.1365-2028.2004.00518.x

255 Henschel, P., Hunter, L. T. B., Coad, L., Abernethy, K. A., & Mühlenberg, M. (2011).
 256 Leopard prey choice in the Congo Basin rainforest suggests exploitative competition with
 257 human bushmeat hunters. *Journal of Zoology*, 285, 11-20. doi:10.1111/j.1469-
 258 7998.2011.00826.x

259 Jacobson, A. P., Gerngross, P., Lemeris Jr, J. R., Schoonover, R. F., Anco, C., Breitenmoser-
 260 Würsten, C., . . . Dollar, L. (2016). Leopard (*Panthera pardus*) status, distribution, and the
 261 research efforts across its range. *PeerJ*, 4, e1974. doi:10.7717/peerj.1974

262 Kabanda, T. A. (2003). Climate. In K. Berger, J. E. Crafford, I. Gaigher, M. J. Gaigher, N.
 263 Hahn, & I. Macdonald (Eds.), *A first synthesis of the environmental, biological and cultural*
 264 *assets of the Soutpansberg* (pp. 10-11). Louis Trichardt: Leach Printers & Signs.

265 Karanth, K. U., & Sunquist, M. E. (1995). Prey selection by tiger, leopard and dhole in
 266 tropical forests. *Journal of Animal Ecology*, 64, 439-450. doi:10.2307/5647

267 Karanth, K. U., & Sunquist, M. E. (2000). Behavioural correlates of predation by tiger
 268 (*Panthera tigris*), leopard (*Panthera pardus*) and dhole (*Cuon alpinus*) in Nagarahole, India.
 269 *Journal of Zoology*, 250, 255-265. doi:10.1111/j.1469-7998.2000.tb01076.x

270 Kent, V. T., & Hill, R. A. (2013). The importance of farmland for the conservation of brown
 271 hyaena, *Parahyaena brunnea*. *Oryx*, 47, 431-440. doi:10.1017/S0030605312001007

272 Keogh, H. J. (1983). A photographic reference system of the microstructure of the hair of
 273 southern African bovids. *South African Journal of Wildlife Research*, 13, 89-131.

274 Kinnaird, M. F., & O'Brien, T. G. (2012). Effects of private- land use, livestock
 275 management, and human tolerance on diversity, distribution, and abundance of large African
 276 mammals. *Conservation Biology*, 26, 1026-1039. doi:10.1111/j.1523-1739.2012.01942.x

277 Knott, P., Knott, H., Kruger, J., & Van der Waal, C. (2003). Private game reserves and state
 278 reserves. In K. Berger, J. E. Crafford, I. Gaigher, M. J. Gaigher, N. Hahn, & I. Macdonald
 279 (Eds.), *A first synthesis of the environmental, biological and cultural assets of the*
 280 *Soutpansberg* (pp. 59-60). Louis Trichardt: Leach Printers & Signs.

281 Kuhn, B. F., Wiesel, I., & Skinner, J. D. (2008). Diet of brown hyaenas (*Parahyaena*
 282 *brunnea*) on the Namibian coast. *Transactions of the Royal Society of South Africa*, 63, 150-
 283 159. doi:10.1080/00359190809519219

284 Li, S., McShea, W. J., Wang, D., Shao, L., & Shi, X. (2010). The use of infrared- triggered
 285 cameras for surveying phasianids in Sichuan Province, China. *Ibis*, 152, 299-309.
 286 doi:10.1111/j.1474-919X.2009.00989.x

287 Macdonald, I. A. W., Gaigher, I., Gaigher, R., & Berger, K. (2003). Executive summary. In
 288 K. Berger, J. E. Crafford, I. Gaigher, M. J. Gaigher, N. Hahn, & I. Macdonald (Eds.), *A first*
 289 *synthesis of the environmental, biological and cultural assets of the Soutpansberg* (pp. 4-6).
 290 Louis Trichardt: Leach Printers & Signs.

291 Martins, Q., Horsnell, W. G. C., Titus, W., Rautenbach, T., & Harris, S. (2011). Diet
 292 determination of the Cape Mountain leopards using global positioning system location
 293 clusters and scat analysis. *Journal of Zoology*, 283, 81-87. doi:10.1111/j.1469-
 294 7998.2010.00757.x

295 Maude, G., & Mills, M. G. L. (2005). The comparative feeding ecology of the brown hyaena
 296 in a cattle area and a national park in Botswana. *South African Journal of Wildlife Research*,
 297 35, 201-214.

298 Mills, M. G. L. (1984). The comparative behavioural ecology of the brown hyaena *Hyaena*
 299 *brunnea* and the spotted hyaena *Crocuta crocuta* in the southern Kalahari. *Koedoe*, 27, 237-
 300 247. doi:10.4102/koedoe.v27i2.583

301 Mills, M. G. L. (2015). Living near the edge: a review of the ecological relationships between
 302 large carnivores in the arid Kalahari. *South African Journal of Wildlife Research*, 45, 127-
 303 137. doi:10.3957/056.045.0127

304 Mills, M. G. L., & Mills, M. E. J. (1978). The diet of the brown hyaena *Hyaena brunnea* in
 305 the southern Kalahari. *Koedoe*, 21, 125-149. doi:10.4102/koedoe.v21i1.968

306 Mori, E., Maggini, I., & Menchetti, M. (2014). When quills kill: the defense strategy of the
 307 crested porcupine *Hystrix cristata* L., 1758. *Mammalia*, 78, 229-234. doi:10.1515/mammalia-
 308 2013-0126

309 Navia, A. F., Mejía-Falla, P. A., & Giraldo, A. (2007). Feeding ecology of elasmobranch
 310 fishes in coastal waters of the Colombian Eastern Tropical Pacific. *BMC Ecology*, 7, 1-10.
 311 doi:10.1186/1472-6785-7-8

312 Negrões, N., Sarmiento, P., Cruz, J., Eira, C., Revilla, E., Fonseca, C., . . . Silveira, L. (2010).
 313 Use of camera-trapping to estimate puma density and influencing factors in central Brazil.
 314 *Journal of Wildlife Management*, 74, 1195-1203. doi:10.2193/2009-256

315 Nilsen, E. B., Christianson, D., Gaillard, J., Halley, D., Linnell, J. D. C., Odden, M., . . .
 316 Zimmermann, B. (2012). Describing food habits and predation: field methods and statistical
 317 considerations. In L. Boitani & R. A. Powell (Eds.), *Carnivore ecology and conservation: a*
 318 *handbook of techniques* (pp. 256-272). Oxford: Oxford University Press.

319 O'Brien, T. G., Kinnaird, M. F., & Wibisono, H. T. (2003). Crouching tigers, hidden prey:
 320 Sumatran tiger and prey populations in a tropical forest landscape. *Animal Conservation*, 6,
 321 131-139. doi:10.1017/s1367943002003172

322 Owens, M. J., & Owens, D. D. (1978). Feeding ecology and its influence on social
 323 organization in brown hyenas (*Hyaena brunnea*, Thunberg) of the central Kalahari desert.
 324 *East African Wildlife Journal*, 16, 113-135. doi:10.1111/j.1365-2028.1978.tb00433.x

325 Pianka, E. R. (1973). The structure of lizard communities. *Annual Review of Ecology and*
 326 *Systematics*, 4, 53-74. doi:10.1146/annurev.es.04.110173.000413

327 Quinn, G. P., & Keough, M. J. (2002). *Experimental designs and data analysis for biologists*.
328 Cambridge: Cambridge University Press.

329 R Development Core Team. (2017). R: a language and environment for statistical computing,
330 [Computer Program]. Vienna: The R Foundation for Statistical Computing. Retrieved from
331 <https://www.r-project.org/>

332 Rovero, F., & Marshall, A. R. (2009). Camera trapping photographic rate as an index of
333 density in forest ungulates. *Journal of Applied Ecology*, 46, 1011-1017. doi:10.1111/j.1365-
334 2664.2009.01705.x

335 Seiler, N. (2010). SEM-Atlas of hair structures of South African mammals. *Mammalia*, 74,
336 281-290. doi:10.1515/mamm.2010.033

337 Slater, K., & Muller, K. (2014). The diet of brown hyaenas (*Hyaena brunnea*) in Shamwari
338 Game Reserve, Eastern Cape, South Africa. *Koedoe*, 56, 1-5. doi:10.4102/koedoe.v56i1.143

339 Stein, A. B., Fuller, T. K., & Marker, L. L. (2013). Brown hyaena feeding ecology on
340 Namibian farmlands. *South African Journal of Wildlife Research*, 43, 27-32.
341 doi:10.3957/056.043.0115

342 Stuart, C., & Stuart, T. (2003). *A field guide to the tracks and signs of southern and east*
343 *African wildlife* (Third ed.). Cape Town: Struik.

344 Taru, P., & Backwell, L. (2013). Identification of fossil hairs in *Parahyaena brunnea*
345 coprolites from Middle Pleistocene deposits at Gladysvale Cave, South Africa. *Journal Of*
346 *Archaeological Science*, 40, 3674-3685. doi:10.1016/j.jas.2013.04.031

347 van der Merwe, I., Tambling, C. J., Thorn, M., Scott, D. M., Yarnell, R. W., Green, M., . . .
 348 Bateman, P. W. (2009). An assessment of diet overlap of two mesocarnivores in the North
 349 West Province, South Africa. *African Zoology*, 44, 288-291. doi:10.3377/004.044.0217
 350 Williams, K. S. (2017). *Human-brown hyaena relationships and the role of mountainous*
 351 *environments as refuges in a postcolonial landscape*. (PhD), Durham University, Durham.
 352 Williams, S. T., Williams, K. S., Lewis, B. P., & Hill, R. A. (2017). Population dynamics and
 353 threats to an apex predator outside of protected areas: Implications for carnivore
 354 management. *Royal Society Open Science*, 4. doi:10.1098/rsos.161090
 355 Yarnell, R. W., Phipps, W. L., Burgess, L. P., Ellis, J. A., Harrison, S. W. R., Dell, S., . . .
 356 Scott, D. M. (2013). The influence of large predators on the feeding ecology of two African
 357 mesocarnivores: the black-backed jackal and the brown hyaena. *South African Journal of*
 358 *Wildlife Research*, 43, 155-166. doi:10.3957/056.043.0206

359

360

361 **Table 1** Occurrence of mammalian prey species identified in brown hyaena and leopard scats collected in the
362 western Soutpansberg Mountains, South Africa, between July 2011 and December 2015. Prey size groupings are
363 based on classifications by Mills and Mills (1978).

Prey species	Brown hyaena (n=137)		Leopard (n=237)	
	Corrected Occurrences	Corrected frequency of occurrence %	Corrected Occurrences	Corrected frequency of occurrence %
Large mammals (> 50 kg)				
Blesbok, <i>Damaliscus pygargus</i>	1	0.73		
Blue wildebeest, <i>Connochaetes taurinus</i>			0.33	0.14
Bushpig, <i>Potamochoerus larvatus</i>	7.58	5.53	24	10.13
Common warthog, <i>Phacochoerus africanus</i>	9.17	6.69	6.33	2.67
Gemsbok, <i>Oryx gazella</i>	1	0.73		
Giraffe, <i>Giraffa camelopardalis</i>	1	0.73		
Greater kudu, <i>Tragelaphus strepsiceros</i>	8.08	5.9	6	2.53
Nyala, <i>Tragelaphus angasii</i>	5	3.65		
Sable, <i>Hippotragus niger</i>	0.33	0.24		
Waterbuck, <i>Kobus ellipsiprymnus</i>	1.83	1.34	4	1.69
Zebra, <i>Equus quagga</i>	1.75	1.28		
Domestic livestock				
Cow, <i>Bos taurus</i>	5.08	3.71		
Goat, <i>Capra aegagrus hircus</i>	4.33	3.16		
Sheep, <i>Ovis aries</i>	0.5	0.36		
Medium mammals (16 - 50 kg)				
Aardvark, <i>Orycteropus afer</i>	1.33	0.97		
Brown hyaena, <i>Hyaena brunnea</i>	1.17	0.85		
Bushbuck, <i>Tragelaphus scriptus</i>	20	14.6	80.83	34.11
Chacma baboon, <i>Papio ursinus</i>	11.83	8.64	16.83	7.1
Common duiker, <i>Sylvicapra grimmia</i>	10.25	7.48	15.67	6.61
Domestic dog, <i>Canis lupus familiaris</i>	0.5	0.36		
Grey rhebok, <i>Pelea capreolus</i>			1	0.42
Impala, <i>Aepyceros melampus</i>	8.08	5.9	3.83	1.62
Mountain reedbuck, <i>Redunca fulvorufula</i>			1.5	0.63
Small mammals (1 – 15 kg)				
African civet, <i>Civettictis civetta</i>	0.33	0.24		
Black-backed jackal, <i>Canis mesomelas</i>	1	0.73		
Cape porcupine, <i>Hystrix africaeaustralis</i>	0.67	0.49	5.17	2.18
Gambian giant rat, <i>Cricetomys gambianus</i>	1.25	0.91	1	0.42
Klipspringer, <i>Oreotragus oreotragus</i>			5.67	2.39
Large spotted genet, <i>Genetta maculata</i>	0.5	0.36		
Mongoose, Family: Herpestidae [†]	2.5	1.82		
Red duiker, <i>Cephalophus natalensis</i>	9.17	6.69	8.83	3.73
Rock dassie, <i>Procavia capensis</i>	1.33	0.97	12.17	5.13
Samango monkey, <i>Cercopithecus albogularis</i>	3.83	2.8	7	2.95

Sharpe's grysbok, <i>Raphicerus sharpei</i>	1	0.73		
Steenbok, <i>Raphicerus campestris</i>	0.5	0.36		
Thick-Tailed Bushbaby, <i>Otolemur crassicaudatus</i>			1	0.42
Vervet monkey, <i>Chlorocebus pygerythrus</i>	5.17	3.77	19.83	8.37
Yellow spotted dassie, <i>Heterohyrax brucei</i>	4.08	2.98	11.5	4.85
Very small mammals (< 1 kg)				
Four striped mouse, <i>Rhabdomys pumilio</i>	1.83	1.34	1	0.42
House rat, <i>Rattus rattus</i>	0.92	0.67		
Lesser bushbaby, <i>Galago moholi</i>	1	0.73		
Lesser red musk shrew, <i>Crocidura hirta</i>	0.75	0.55		
Namaqua rock mouse, <i>Aethomys namaquensis</i>			0.5	0.21
Rock elephant shrew, <i>Elephantulus myurus</i>	0.5	0.36	2	0.84
Swamp musk shrew, <i>Crocidura mariquensis</i>	0.83	0.61		
Woodland dormouse, <i>Graphiurus murinus</i>			0.5	0.21
Total		100		100

[†] It was possible to identify mongooses only to a Family level.

Fig. 1 Locations of brown hyaena and leopard scats collected, and minimum convex polygon (MCP) of the area covered by camera traps in the Soutpansberg Mountains.




Fig. 2 (a) Corrected frequency of occurrence (CFO) of prey species in brown hyaena scats and corrected frequency of occurrence of prey species in leopard scats. (b) Corrected frequency of occurrence (CFO) of prey species in brown hyaena scats and relative abundance index (RAI) of prey species. Shaded area represents 95% confidence interval.

